



SA EQUINE
HEALTH & PROTOCOLS
EXPORTS SOUTH AFRICA



African horse sickness control

Surveillance report

**Sentinel surveillance
2023**

**JD Grewar, CT Weyer & LS van Helden
2023**



**Western Cape
Government**
FOR YOU

Overview

The African horse sickness (AHS) sentinel surveillance program provides additional confidence of AHS freedom in the AHS free (FZ) and surveillance zones (SZ) of South Africa. The program incorporates the monthly sampling of recruited horses proportionately selected within the zones based on the estimated underlying population. Historically the program had two components – a sero-sentinel program that evaluated the changing serological status of horses on a month-to-month basis; and a PCR-based program that is used to detect the presence of AHS viral RNA within recruits. In 2023 the serological component of the program was suspended with the PCR-based program remaining the focus of the program for the foreseeable future.

The PCR sampling target is drawn up to detect AHS at approximately a 2% minimum expected prevalence (with a 95% confidence level), resulting in a monthly sampling target of 150 animals. The vaccination status of PCR sentinels does not influence their recruitment, unless vaccination against AHS took place sufficiently recently to result in positive PCR results on initial testing.

A detailed description of the original program is available in the [January 2016 Western Cape Epidemiology Report](#). The summary report for the 2021/22 season can be found in the [July 2023 Epidemiology Report](#). All other reports can be found at www.myhorse.org.za.

Viral RNA PCR testing is generally performed at the Stellenbosch Provincial Veterinary Laboratory (SPVL) in the sentinel surveillance program. During 2023 however, samples were tested at the University of Pretoria/Equine Research Center's Molecular Diagnostics Laboratory as the SPVL was undergoing renovations. The PCR test method used is a University of Pretoria (Equine Research Center) developed and WOAHP validated real-time RT-PCR (Guthrie et al. 2013).

This report covers the AHS sentinel program for the 2023 calendar year. The results confirm that it is unlikely that AHS was circulating in the AHS free and surveillance zone during this period.

General overview of sampling and results

1719 PCR sentinel samples were analysed from 63 different farms at an average of 143 samples from, on average, 48 different farms per month. All samples tested negative.

Investigations

There were no follow-up investigations as a result of suspect or positive laboratory results during the year. There was however an investigation into a fever reaction on a sentinel property (holding ID 79 with 8 resident sentinels – see Figure 4 and Figure 5) in December 2023, however the fever was not in a sentinel animal. The affected animal presented with fever and mild ataxia on the 22 December 2023. Infection with Equine encephalosis virus (EEV) was considered a differential diagnosis by the attending veterinarian.

The sample that was taken tested negative for both AHS and EEV RNA (Lab ref E240010). There were 4 sentinel farms in total, including farm 79, within a 10 km radius of the fever case. A total of 16

sentinels from within this area were tested (all negative) in the December 2023 sentinel sampling cohort.

Spatial considerations

The sentinel surveillance program is based on a proportional sampling system with most sentinels in areas of the surveillance area that have the highest population of horses. **Error! Reference source not found.** and **Error! Reference source not found.** show the underlying population and current sentinel farms and the monthly average distribution of sentinels in the PCR sentinel program. Some improvement has been made in the Paarl area when compared to 2021/2022, with the Mitchells Plain area still requiring most improvement.




Figure 1: The underlying population of horses in the Surveillance and Free Zones of South Africa. These populations have been revised based on new population data collected between 1 April 2016 and December 2023.




Figure 2: A map showing the AHS surveillance and free zone where PCR-sentinel surveillance has taken place during 2023. The map depicts the various areas with their target PCR samples to detect a 2% minimum expected prevalence using a proportional sampling frame. The orange areas are areas where PCR-sentinels were lacking on average while the light green to green areas show where surplus PCR-sentinels were sampled. Cream areas depict where the target was generally attained.

Surveillance system evaluation

The surveillance program is designed to detect AHS in the AHS surveillance zone at a minimum expected prevalence of 2% (MEP). In this section of the report, we establish the monthly sensitivity of the surveillance program. Note that previous analyses evaluated the program at a 5% MEP based on EU 2008/698 requirements – this legislation is now repealed and, since the program aims at a 2% MEP, the evaluation thereof has been adjusted to this level.

Parameters used in this evaluation are shown in Table 1 and analysis is based on evaluating sensitivity of surveillance programs (Martin et al. 2007). The historical surveillance outcome is considered as it provides information that aids in determining an accurate final probability of freedom as of December

2023. The final probability of freedom from Sept 2016 through December 2023 (88 months) was 88.8% - see Figure 3).

The sensitivity of the sentinel surveillance stays around 25%-30%. This is the seventh AHS season running where cases of the disease have not been detected in the AHS surveillance and free area, although an outbreak of AHS occurred in the AHS protection zone in 2021

Parameter	Value	Comments
<i>p</i> Intro	0.03	During periods where not outbreaks in the AHS controlled area are present. Based on historical outbreaks in the region.
	0.3	During periods where outbreaks are present in the AHS controlled area – estimate made increasing probability of introduction 10X the normal rate
Population at risk – total herds	1478	Data captured between 1 April 2016 and Dec 2023 for the AHS surveillance and free zones.
Sentinel farm populations	Various	Based on herd size as of Dec 2023 . The assumption is made that herd size would not change substantially on the sentinel properties over the period reviewed.
Sentinels tested per herd per surveillance period	Various	Actual tested data
Unit design prevalence (P_A^*)	0.02	Design prevalence at animal level
Herd design prevalence (P_H^*)	0.02	Design prevalence at herd level based on prior outbreaks (median value taken) in the controlled area assuming a herd PAR of the zones affected by each outbreak.
Test sensitivity	0.978	As published (Guthrie et al. 2013). Note that while serology was taken into consideration, for this analysis all horses that were tested on serology were tested on PCR – hence the use of a single test sensitivity across the analysis
Initial Prior confidence of Freedom	0.5	Note that when evaluating the season independently the prior of 0.5 is used in the first surveillance period (September 2021). When evaluating the past 6 years between Sept 2016 and Dec 2022 the initial prior is 0.5 but relates to September 2016.

Table 1: Parameters used to establish sentinel system probability and sensitivity of freedom for African horse sickness




Figure 3: The sentinel surveillance sensitivity of individual surveillance periods (*SeP* - dots) with probability of freedom curve (red line) for the past seven surveillance seasons: the season currently reviewed is the 2023 calendar year. Probability of AHS introduction of 3% is set for periods where no AHS outbreaks are present in the AHS controlled area (grey line at 0.03 on y-axis) but at 10X that rate for where outbreaks are present as in April and May 2021 in the Cederberg AHS Protection zone.

Discussion and Conclusion

The primary goal of demonstrating AHS freedom 2023 was achieved. A 7-year review of sentinel results show that the probability of freedom attained for this program, at an animal design prevalence of 2% and herd-level design prevalence of 2%, shows a 88.8% probability of freedom from AHS in the AHS surveillance and free zones. This level was achieved in the face of the AHS outbreak that occurred ~ 88km from the border of the AHS surveillance zone in 2021. It further does not take into consideration the passive surveillance component. Spatial representativeness remains challenging, but gains have been made in the Paarl region.

References and acknowledgements

This program would not be possible without the support of the horse owners in the AHS surveillance zone who freely give of their time and resources to allow and facilitate the monthly sampling of horses. We are grateful to the University of Pretoria Molecular Genetics Laboratory who performed the testing of samples this season.

In this season the sentinel program was again achieved through collaboration between the Western Cape Department of Agriculture (Veterinary Services) and SAEHP. In this regard we specifically acknowledge Dr Tasneem Anthony. The WCDOA also currently fund the testing costs associated with the program. We are grateful to the SAEHP team who are directly involved with the program: Esthea Russouw and Lize Germishuys.

Software and systems references

Evan Sergeant (2016). RSurveillance: Design and Analysis of Disease Surveillance Activities. R package version 0.2.0

H. Wickham (2009). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

Hadley Wickham (2011). The Split-Apply-Combine Strategy for Data Analysis. Journal of Statistical Software, 40(1), 1-29.

Hadley Wickham (2016). scales: Scale Functions for Visualization. R package version 0.4.0.

Achim Zeileis and Gabor Grothendieck (2005). zoo: S3 Infrastructure for Regular and Irregular Time Series. Journal of Statistical Software, 14(6), 1-27

Joe Conway, Dirk Eddelbuettel, Tomoaki Nishiyama, Sameer Kumar Prayaga and Neil Tiffin (2016). RPostgreSQL: R interface to the PostgreSQL database system. R package version 0.4-1.

Literature references

Guthrie, A.J. et al., 2013. Diagnostic accuracy of a duplex real-time reverse transcription quantitative PCR assay for detection of African horse sickness virus. Journal of Virological Methods, 189(1), pp.30–35.

Martin, P.A.J., Cameron, A.R. & Greiner, M., 2007. Demonstrating freedom from disease using multiple complex data sources. 1: A new methodology based on scenario trees. Preventive Veterinary Medicine, 79(2–4), pp.71–97

Appendix – Investigation images




Figure 4: Holding 79 in relation to AHS controlled area and all other sentinel properties depicted by black dots




Figure 5: Close up of the 5 and 10 km buffers surrounding the farm investigated from Figure 5, showing other sentinel farms with their proportional number of sentinels resident